


### AMENDMENTS TO THE CLAIMS

Kindly amend claims 1 – 4 and cancel claims 8 – 9 without prejudice to the subject matter involved.  
This listing of claims will replace all prior versions, and listings, of claims in the application:

#### Listing of Claims:

1. (Currently amended) A compound of formula



wherein

A<sub>0</sub>, A<sub>1</sub> and A<sub>2</sub> are each independently of the others a bond or a C<sub>1</sub>-C<sub>6</sub>alkylene bridge which is unsubstituted or substituted by from one to six identical or different substituents selected from halogen and C<sub>3</sub>-C<sub>8</sub>cycloalkyl;

A<sub>3</sub> is a C<sub>1</sub>-C<sub>6</sub>alkylene bridge which is unsubstituted or substituted by from one to six identical or different substituents selected from halogen and C<sub>3</sub>-C<sub>8</sub>cycloalkyl;

Y is O, NR<sub>11</sub>, S, SO or SO<sub>2</sub>;

M is O or NOR<sub>6</sub>,

X<sub>1</sub> and X<sub>2</sub> are each independently of the other fluorine, chlorine or bromine;

R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> are each independently of the others H, halogen, OH, SH, CN, nitro, C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>haloalkyl, C<sub>1</sub>-C<sub>6</sub>alkylcarbonyl, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>2</sub>-C<sub>6</sub>haloalkenyl, C<sub>2</sub>-C<sub>6</sub>alkynyl, C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>1</sub>-C<sub>6</sub>haloalkoxy, C<sub>2</sub>-C<sub>6</sub>alkenyloxy, C<sub>2</sub>-C<sub>6</sub>haloalkenyloxy, C<sub>2</sub>-C<sub>6</sub>alkynyloxy, -S(=O)-C<sub>1</sub>-C<sub>6</sub>alkyl, -S(=O)<sub>2</sub>-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>alkoxycarbonyl or C<sub>3</sub>-C<sub>6</sub>haloalkynyloxy; the substituents R<sub>3</sub> being independent of one another when m is 2;

Q is O, NR<sub>11</sub>, S, SO or SO<sub>2</sub>;

W is O, NR<sub>11</sub>, S, SO, SO<sub>2</sub>, -C(=O)-O-, -O-C(=O)-, -C(=O)-NR<sub>11</sub>- or -NR<sub>11</sub>-C(=O)-;

T is a bond, O, NR<sub>11</sub>, S, SO, SO<sub>2</sub>, -C(=O)-O-, -O-C(=O)-, -C(=O)-NR<sub>11</sub>- or -NR<sub>11</sub>-C(=O)-;

D is CH or N;

$R_4$  is H, halogen, OH, SH, CN, nitro,  $C_1$ - $C_6$ alkyl,  $C_1$ - $C_6$ haloalkyl,  $C_1$ - $C_6$ alkylcarbonyl,  $C_2$ - $C_6$ alkenyl,  $C_2$ - $C_6$ haloalkenyl,  $C_2$ - $C_6$ alkynyl,  $C_1$ - $C_6$ alkoxy,  $C_1$ - $C_6$ haloalkoxy,  $C_2$ - $C_6$ alkenyloxy,  $C_2$ - $C_6$ haloalkenyloxy,  $C_2$ - $C_6$ alkynyloxy,  $-S(=O)-C_1-C_6$ alkyl,  $-S(=O)_2-C_1-C_6$ alkyl,  $C_1$ - $C_6$ alkoxycarbonyl,  $C_3$ - $C_6$ haloalkynyloxy,  $NH_2$ ,  $NH(C_1-C_6$ alkyl) or  $N(C_1-C_6$ alkyl) $_2$  wherein the two alkyl groups are independent of one another; the substituents  $R_4$  being independent of one another when k is greater than 1;

$R_5$  is  $C_1$ - $C_{12}$ alkoxy- $C_1$ - $C_{12}$ alkyl or heterocyclyl;  $C_1$ - $C_{12}$ alkyl substituted by from one to five substituents selected from the group consisting of  $-N_3$ ,  $NO_{27}$ , OH,  $C_3$ - $C_6$ cycloalkyl,  $C_3$ - $C_6$ cycloalkoxy,  $C_4$ - $C_6$ alkoxy,  $C_4$ - $C_6$ haloalkoxy,  $C_2$ - $C_6$ alkenyloxy,  $C_2$ - $C_6$ haloalkenyloxy,  $C_3$ - $C_6$ alkynyloxy,  $C_3$ - $C_6$ haloalkynyl,  $C_3$ - $C_6$ haloalkynyloxy,  $C_3$ - $C_6$ cycloalkyl- $C_4$ - $C_6$ alkoxy,  $C_4$ - $C_6$ alkylcarbonyl,  $C_4$ - $C_6$ haloalkylcarbonyl,  $C_1$ - $C_6$ alkoxy- $C_4$ - $C_6$ alkoxy,  $-P(=O)(OC_1-C_6$ alkyl) $_{27}$ ,  $-S(O)_q-R_{143}$ ,  $NH_{27}$ ,  $NH(C_1-C_6$ alkyl),  $N(C_1-C_6$ alkyl) $_2$ , wherein the two alkyl groups are independent of one another,  $-N(R_{27})_2$  wherein the two  $R_{27}$ s are independent of one another and  $-NR_{144}S(O)_qR_{146}$ ;

$C_3$ - $C_6$ cycloalkyl substituted by from one to five identical or different substituents selected from the group consisting of  $C_4$ - $C_6$ alkyl, halogen, CN,  $NO_{27}$ , OH,  $C_4$ - $C_6$ alkoxy,  $C_4$ - $C_6$ haloalkoxy,  $NH_{27}$ ,  $NH(C_1-C_6$ alkyl) and  $N(C_1-C_6$ alkyl) $_2$  wherein the two alkyl groups are independent of one another;

$-N(R_{27})_2$  wherein the two  $R_{27}$ s are independent of one another;

$-C(=O)-O-R_{61}-C(=O)-R_{61}-C(=O)-NH-R_{61}-C(=N-O-R_{61})R_{1401}-C(=N-NH-R_{61})R_{1401}-C_2-C_6$ alkenyl;  $C_2$ - $C_6$ alkynyl; heterocyclyl; or

$-NR_{144}S(O)_qR_{146}$

wherein the alkenyl and alkynyl radicals are unsubstituted or, depending upon the possibilities of substitution, substituted by from one to five identical or different substituents selected from the group consisting of halogen,  $-N_3$ , CN,  $NO_{27}$ , OH,  $C_3$ - $C_6$ cycloalkyl,  $C_4$ - $C_6$ alkoxy,  $C_4$ - $C_6$ haloalkoxy,  $C_2$ - $C_6$ alkenyloxy,  $C_2$ - $C_6$ haloalkenyloxy,  $C_3$ - $C_6$ alkynyloxy,  $C_3$ - $C_6$ haloalkynyloxy,  $C_3$ - $C_6$ cycloalkyl- $C_4$ - $C_6$ alkoxy,  $C_4$ - $C_6$ alkylcarbonyl,  $C_4$ - $C_6$ haloalkylcarbonyl,  $C_1$ - $C_6$ alkoxycarbonyl,  $C_4$ - $C_6$ alkylcarbonyl- $C_4$ - $C_6$ alkyl,  $C_1$ - $C_6$ alkoxycarbonyl- $C_1$ - $C_6$ alkyl,  $C_3$ - $C_6$ haloalkynyl,  $C_4$ - $C_6$ alkoxy- $C_1$ - $C_6$ alkyl,  $C_1$ - $C_6$ haloalkoxy- $C_1$ - $C_6$ alkyl,  $C_2$ - $C_6$ alkenyloxy- $C_4$ - $C_6$ alkyl,  $C_2$ - $C_6$ haloalkenyloxy- $C_1$ - $C_6$ alkyl,  $C_3$ - $C_6$ alkynyloxy- $C_1$ - $C_6$ alkyl,  $-P(=O)(OC_1-C_6$ alkyl) $_{27}$ ,  $-S(O)_q-R_{143}$ ,  $NH_{27}$ ,  $NH(C_1-C_6$ alkyl) and  $N(C_1-C_6$ alkyl) $_{27}$  wherein the two alkyl groups are independent of one another;

and wherein the heterocyclyl radical mentioned under  $R_5$  are unsubstituted or, depending upon the possibilities of substitution, substituted by from one to five substituents selected from

halogen, CN, NO<sub>2</sub>, OH, SH, C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>haloalkyl, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>2</sub>-C<sub>6</sub>haloalkenyl, C<sub>3</sub>-C<sub>6</sub>alkynyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>1</sub>-C<sub>6</sub>haloalkoxy, C<sub>2</sub>-C<sub>6</sub>alkenyloxy, C<sub>2</sub>-C<sub>6</sub>haloalkenyloxy, C<sub>3</sub>-C<sub>6</sub>alkynyloxy, C<sub>3</sub>-C<sub>6</sub>haloalkynyloxy, C<sub>3</sub>-C<sub>6</sub>cycloalkyl-C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>1</sub>-C<sub>6</sub>alkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>haloalkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>alkoxycarbonyl, C<sub>1</sub>-C<sub>6</sub>alkylcarbonyl-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>alkoxycarbonyl-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>alkylthio, C<sub>2</sub>-C<sub>6</sub>alkenylthio, C<sub>3</sub>-C<sub>6</sub>alkynylthio, C<sub>3</sub>-C<sub>6</sub>cycloalkyl-C<sub>1</sub>-C<sub>6</sub>alkylthio, C<sub>3</sub>-C<sub>6</sub>haloalkynyl, C<sub>2</sub>-C<sub>6</sub>haloalkenylthio, C<sub>1</sub>-C<sub>6</sub>haloalkylthio, C<sub>1</sub>-C<sub>6</sub>alkoxy-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>haloalkoxy-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>2</sub>-C<sub>6</sub>alkenyloxy-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>2</sub>-C<sub>6</sub>haloalkenyloxy-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>3</sub>-C<sub>6</sub>alkynyloxy-C<sub>1</sub>-C<sub>6</sub>alkyl, NH<sub>2</sub>, NH(C<sub>1</sub>-C<sub>6</sub>alkyl), N(C<sub>1</sub>-C<sub>6</sub>alkyl)<sub>2</sub> wherein the two alkyl groups are independent of one another, C<sub>1</sub>-C<sub>6</sub>alkylcarbonylamino, C<sub>1</sub>-C<sub>6</sub>haloalkylcarbonylamino, C<sub>1</sub>-C<sub>6</sub>alkoxycarbonylamino and C<sub>1</sub>-C<sub>6</sub>alkylaminocarbonylamino;

or, when A<sub>6</sub> is a C<sub>4</sub>-C<sub>6</sub>alkylene bridge, R<sub>6</sub> is C<sub>2</sub>-C<sub>6</sub>alkylene bonded to one of the carbon atoms of A<sub>6</sub>;

or, when R<sub>4</sub> and a group C(=NOR<sub>6</sub>)R<sub>6</sub> are in the ortho position relative to one another, R<sub>4</sub> and R<sub>6</sub> together form a C<sub>2</sub>-C<sub>6</sub>alkylene bridge wherein one or two CH<sub>2</sub> groups each independently of the other may be replaced by O, NR<sub>12</sub>, S or SO, and wherein the CH<sub>2</sub> groups are unsubstituted or mono- or di-substituted by halogen, OH, SH, CN, nitro, C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>haloalkyl, C<sub>1</sub>-C<sub>6</sub>alkoxy or C<sub>1</sub>-C<sub>6</sub>haloalkoxy;

R<sub>6</sub> is H, C<sub>1</sub>-C<sub>12</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>cycloalkyl, C<sub>1</sub>-C<sub>6</sub>alkylcarbonyl, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>2</sub>-C<sub>6</sub>alkynyl, aryl, heterocycl or benzyl, wherein the alkyl, cycloalkyl, alkenyl and alkynyl radicals are unsubstituted or, depending upon the possibilities of substitution, substituted by from one to five identical or different substituents selected from the group consisting of halogen, -N<sub>3</sub>, CN, NO<sub>2</sub>, OH, SH, C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>1</sub>-C<sub>6</sub>haloalkoxy, C<sub>2</sub>-C<sub>6</sub>alkenyloxy, C<sub>2</sub>-C<sub>6</sub>haloalkenyloxy, C<sub>3</sub>-C<sub>6</sub>alkynyloxy, C<sub>3</sub>-C<sub>6</sub>haloalkynyloxy, C<sub>3</sub>-C<sub>6</sub>cycloalkyl-C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>1</sub>-C<sub>6</sub>alkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>haloalkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>alkoxycarbonyl, C<sub>1</sub>-C<sub>6</sub>alkylcarbonyl-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>alkoxycarbonyl-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>alkylthio, C<sub>2</sub>-C<sub>6</sub>alkenylthio, C<sub>3</sub>-C<sub>6</sub>alkynylthio, C<sub>3</sub>-C<sub>6</sub>cycloalkyl-C<sub>1</sub>-C<sub>6</sub>alkylthio, C<sub>3</sub>-C<sub>6</sub>haloalkynyl, C<sub>2</sub>-C<sub>6</sub>haloalkenylthio, C<sub>1</sub>-C<sub>6</sub>haloalkylthio, C<sub>1</sub>-C<sub>6</sub>alkoxy-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>haloalkoxy-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>2</sub>-C<sub>6</sub>alkenyl-oxy-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>2</sub>-C<sub>6</sub>haloalkenyloxy-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>3</sub>-C<sub>6</sub>alkynyloxy-C<sub>1</sub>-C<sub>6</sub>alkyl, NH<sub>2</sub>, NH(C<sub>1</sub>-C<sub>6</sub>alkyl), N(C<sub>1</sub>-C<sub>6</sub>alkyl)<sub>2</sub> wherein the two alkyl groups are independent of one another, C<sub>1</sub>-C<sub>6</sub>alkyl-carbonylamino, C<sub>1</sub>-C<sub>6</sub>haloalkylcarbonylamino, C<sub>1</sub>-C<sub>6</sub>alkoxycarbonylamino and C<sub>1</sub>-C<sub>6</sub>alkylaminocar-bonylamino;

and the aryl, heterocycl and benzyl radicals are unsubstituted or, depending upon the possibilities of substitution, substituted by from one to five identical or different substituents selected

from the group consisting of halogen, CN, NO<sub>2</sub>, OH, SH, C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>haloalkyl, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>2</sub>-C<sub>6</sub>haloalkenyl, C<sub>3</sub>-C<sub>6</sub>alkynyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>1</sub>-C<sub>6</sub>haloalkoxy, C<sub>2</sub>-C<sub>6</sub>alkenyloxy, C<sub>2</sub>-C<sub>6</sub>haloalkenyloxy, C<sub>3</sub>-C<sub>6</sub>alkynyloxy, C<sub>3</sub>-C<sub>6</sub>haloalkynyloxy, C<sub>3</sub>-C<sub>6</sub>cycloalkyl-C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>1</sub>-C<sub>6</sub>alkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>haloalkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>alkoxycarbonyl, C<sub>1</sub>-C<sub>6</sub>alkylcarbonyl-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>alkoxycarbonyl-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>alkylthio, C<sub>2</sub>-C<sub>6</sub>alkenylthio, C<sub>3</sub>-C<sub>6</sub>alkynylthio, C<sub>3</sub>-C<sub>6</sub>cycloalkyl-C<sub>1</sub>-C<sub>6</sub>alkylthio, C<sub>3</sub>-C<sub>6</sub>haloalkynyl, C<sub>2</sub>-C<sub>6</sub>haloalkenylthio, C<sub>1</sub>-C<sub>6</sub>haloalkylthio, C<sub>1</sub>-C<sub>6</sub>alkoxy-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>haloalkoxy-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>2</sub>-C<sub>6</sub>alkenyloxy-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>2</sub>-C<sub>6</sub>haloalkenyloxy-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>3</sub>-C<sub>6</sub>alkynyloxy-C<sub>1</sub>-C<sub>6</sub>alkyl, NH<sub>2</sub>, NH(C<sub>1</sub>-C<sub>6</sub>alkyl), N(C<sub>1</sub>-C<sub>6</sub>alkyl)<sub>2</sub> wherein the two alkyl groups are independent of one another, C<sub>1</sub>-C<sub>6</sub>alkylcarbonylamino, C<sub>1</sub>-C<sub>6</sub>haloalkylcarbonylamino, C<sub>1</sub>-C<sub>6</sub>alkoxycarbonylamino and C<sub>1</sub>-C<sub>6</sub>alkylaminocarbonylamino;

R<sub>7</sub> is H, C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>3</sub>haloalkyl, C<sub>1</sub>-C<sub>6</sub>alkylcarbonyl, C<sub>1</sub>-C<sub>3</sub>haloalkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>alkoxycarbonyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkylcarbonyl or formyl;

R<sub>8</sub> is H, C<sub>1</sub>-C<sub>12</sub>alkyl substituted by from one to five identical or different substituents selected from halogen, -N<sub>3</sub>, CN, NO<sub>2</sub>, OH, C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>1</sub>-C<sub>6</sub>alkylthio, NH<sub>2</sub>, NH(C<sub>1</sub>-C<sub>6</sub>alkyl), N(C<sub>1</sub>-C<sub>6</sub>alkyl)<sub>2</sub> wherein the two alkyl groups are independent of one another and C<sub>1</sub>-C<sub>6</sub>alkylcarbonylamino; C<sub>3</sub>-C<sub>6</sub>cycloalkyl, C<sub>1</sub>-C<sub>6</sub>alkylcarbonyl, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>2</sub>-C<sub>6</sub>haloalkenyl, C<sub>2</sub>-C<sub>6</sub>alkynyl, C<sub>2</sub>-C<sub>6</sub>haloalkynyl, aryl, heterocyclyl or benzyl, wherein the aryl, heterocyclyl and benzyl radicals are unsubstituted or, depending upon the possibilities of substitution, substituted by from one to five substituents selected from the group consisting of halogen, CN, NO<sub>2</sub>, OH, C<sub>1</sub>-C<sub>6</sub>haloalkyl, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>2</sub>-C<sub>6</sub>haloalkenyl, C<sub>3</sub>-C<sub>6</sub>alkynyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl, C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>1</sub>-C<sub>6</sub>haloalkoxy, C<sub>2</sub>-C<sub>6</sub>alkenyloxy, C<sub>2</sub>-C<sub>6</sub>haloalkenyloxy, C<sub>3</sub>-C<sub>6</sub>alkynyloxy, C<sub>3</sub>-C<sub>6</sub>haloalkynyloxy, C<sub>1</sub>-C<sub>6</sub>alkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>haloalkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>alkoxycarbonyl, C<sub>1</sub>-C<sub>6</sub>alkylthio, C<sub>2</sub>-C<sub>6</sub>alkenylthio, C<sub>3</sub>-C<sub>6</sub>alkynylthio, C<sub>1</sub>-C<sub>6</sub>alkylthio, C<sub>3</sub>-C<sub>6</sub>haloalkynyl, C<sub>1</sub>-C<sub>6</sub>haloalkylthio, C<sub>1</sub>-C<sub>6</sub>alkoxy-C<sub>1</sub>-C<sub>6</sub>alkyl, NH<sub>2</sub>, NH(C<sub>1</sub>-C<sub>6</sub>alkyl), N(C<sub>1</sub>-C<sub>6</sub>alkyl)<sub>2</sub>, C<sub>1</sub>-C<sub>6</sub>alkylcarbonylamino, C<sub>1</sub>-C<sub>6</sub>haloalkylcarbonylamino, C<sub>1</sub>-C<sub>6</sub>alkoxycarbonylamino and C<sub>1</sub>-C<sub>6</sub>alkylaminocarbonylamino;

R<sub>9</sub> is H, C<sub>1</sub>-C<sub>12</sub>alkyl unsubstituted or substituted by from one to five identical or different substituents selected from halogen, CN, NO<sub>2</sub>, OH, C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>1</sub>-C<sub>6</sub>alkylthio, NH<sub>2</sub>, NH(C<sub>1</sub>-C<sub>6</sub>alkyl), N(C<sub>1</sub>-C<sub>6</sub>alkyl)<sub>2</sub> wherein the two alkyl groups are independent of one another and C<sub>1</sub>-C<sub>6</sub>alkylcarbonylamino; C<sub>3</sub>-C<sub>6</sub>cycloalkyl, C<sub>1</sub>-C<sub>6</sub>alkylcarbonyl, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>2</sub>-C<sub>6</sub>haloalkenyl, C<sub>2</sub>-C<sub>6</sub>alkynyl, C<sub>2</sub>-C<sub>6</sub>haloalkynyl, aryl, heterocyclyl or benzyl, wherein the aryl, heterocyclyl and benzyl radicals are unsubstituted or, depending upon the possibilities of substitution, substituted by from one to five substituents selected from the group consisting of halogen, CN, NO<sub>2</sub>, OH,

C<sub>1</sub>-C<sub>6</sub>haloalkyl, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>2</sub>-C<sub>6</sub>haloalkenyl, C<sub>3</sub>-C<sub>6</sub>alkynyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl, C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>1</sub>-C<sub>6</sub>haloalkoxy, C<sub>2</sub>-C<sub>6</sub>alkenyloxy, C<sub>2</sub>-C<sub>6</sub>haloalkenyloxy, C<sub>3</sub>-C<sub>6</sub>alkynyloxy, C<sub>3</sub>-C<sub>6</sub>haloalkynyloxy, C<sub>1</sub>-C<sub>6</sub>alkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>haloalkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>alkoxycarbonyl, C<sub>1</sub>-C<sub>6</sub>alkylthio, C<sub>2</sub>-C<sub>6</sub>alkenylthio, C<sub>3</sub>-C<sub>6</sub>alkynylthio, C<sub>1</sub>-C<sub>6</sub>alkylthio, C<sub>3</sub>-C<sub>6</sub>haloalkynyl, C<sub>1</sub>-C<sub>6</sub>haloalkylthio, C<sub>1</sub>-C<sub>6</sub>alkoxy-C<sub>1</sub>-C<sub>6</sub>alkyl, NH<sub>2</sub>, NH(C<sub>1</sub>-C<sub>6</sub>alkyl), N(C<sub>1</sub>-C<sub>6</sub>alkyl)<sub>2</sub> wherein the two alkyl groups are independent of one another, C<sub>1</sub>-C<sub>6</sub>alkylcarbonylamino, C<sub>1</sub>-C<sub>6</sub>haloalkylcarbonylamino, C<sub>1</sub>-C<sub>6</sub>alkoxycarbonylamino and C<sub>1</sub>-C<sub>6</sub>alkylaminocarbonylamino;

R<sub>10</sub> is H, C<sub>1</sub>-C<sub>12</sub>alkyl unsubstituted or substituted by from one to five identical or different substituents selected from halogen, CN, NO<sub>2</sub>, OH, C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>1</sub>-C<sub>6</sub>alkylthio, NH<sub>2</sub>, NH(C<sub>1</sub>-C<sub>6</sub>alkyl), N(C<sub>1</sub>-C<sub>6</sub>alkyl)<sub>2</sub> and C<sub>1</sub>-C<sub>6</sub>alkylcarbonylamino; C<sub>3</sub>-C<sub>6</sub>cycloalkyl, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>2</sub>-C<sub>6</sub>haloalkenyl, C<sub>2</sub>-C<sub>6</sub>alkynyl, aryl, heterocyclyl or benzyl, wherein the aryl, heterocyclyl and benzyl radicals are unsubstituted or, depending upon the possibilities of substitution, substituted by from one to five identical or different substituents selected from the group consisting of halogen, CN, NO<sub>2</sub>, OH, SH, C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>haloalkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl, C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>1</sub>-C<sub>6</sub>haloalkoxy, C<sub>1</sub>-C<sub>6</sub>alkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>haloalkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>alkoxycarbonyl, C<sub>1</sub>-C<sub>6</sub>alkylcarbonyl-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>alkoxy-C<sub>1</sub>-C<sub>6</sub>alkyl, NH<sub>2</sub>, NH(C<sub>1</sub>-C<sub>6</sub>alkyl), N(C<sub>1</sub>-C<sub>6</sub>alkyl)<sub>2</sub> wherein the two alkyl groups are independent of one another, C<sub>1</sub>-C<sub>6</sub>alkylcarbonylamino, C<sub>1</sub>-C<sub>6</sub>haloalkylcarbonylamino, C<sub>1</sub>-C<sub>6</sub>alkoxycarbonylamino and C<sub>1</sub>-C<sub>6</sub>alkylaminocarbonylamino;

R<sub>11</sub> and R<sub>12</sub> are each independently of the other H, C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>3</sub>haloalkyl, C<sub>1</sub>-C<sub>6</sub>alkylcarbonyl, C<sub>1</sub>-C<sub>3</sub>haloalkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>alkoxycarbonyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl-C<sub>1</sub>-C<sub>6</sub>alkyl or C<sub>3</sub>-C<sub>6</sub>cycloalkylcarbonyl;

R<sub>13</sub> is H, C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>3</sub>-C<sub>6</sub>alkynyl or C<sub>1</sub>-C<sub>6</sub>haloalkyl;

R<sub>14</sub> is H, C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>3</sub>-C<sub>6</sub>alkynyl or C<sub>1</sub>-C<sub>6</sub>haloalkyl;

R<sub>15</sub> is H, C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>3</sub>-C<sub>6</sub>alkynyl or C<sub>1</sub>-C<sub>6</sub>haloalkyl;

k is 0, 1, 2, 3 or 4;

m is 1 or 2; and

q is 0, 1 or 2;

or, where applicable, a possible E/Z isomer, E/Z isomeric mixture and/or tautomer thereof, in each case in free form or in salt form.

2. (Currently amended) The A compound according to claim 1 wherein M is NOR<sub>6</sub>.

3. (Currently amended) The A compound according to claim 1 wherein M is O.
4. (Currently amended) The A compound according to claim 1 in free form.
5. (Previously presented) A compound according to claim 1 wherein X<sub>1</sub> and X<sub>2</sub> are chlorine or bromine.
6. (Previously presented) A compound according to claim 1 wherein D is CH.
7. (Previously presented) A compound according to claim 1 wherein A<sub>3</sub> is straight-chain alkylene bridge.
8. (Canceled)
9. (Canceled)
10. (Previously presented) A pesticidal composition which comprises as active ingredient at least one compound defined in claim 1, in free form or in agrochemically acceptable salt form, and at least one adjuvant.
11. (Original) A method of controlling pests which comprises applying a pesticidal composition as defined in claim 10 to the pests or to the locus thereof.